

Randomized Rounding

Previously

Probability

Events, probabilities,
conditional probabilities,
union bound

Random variables,
expected value,
linearity of expectation
Markov's inequality

Algorithms

Approx. 3-SAT,

Big idea: we only want the average

$$\begin{array}{l} \text{Linearity of Expectation} \\ E[X+Y] = E[X] + E[Y] \quad (\text{stochastic variable}) \\ \text{"average sum = sum of averages"} \end{array}$$

let $Y_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{otherwise} \end{cases} \quad (i=1, \dots, m)$

$$E[Y_i] = 7/8 \quad (1 \text{ clause case})$$

$$E[\text{\# clauses satisfied}] = E\left[\sum_{i=1}^m Y_i\right] = \sum_{i=1}^m E[Y_i] = 7/8 m$$

let $Y_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{otherwise} \end{cases}$

$$Z = \sum_{i=1}^m Y_i = \text{\# clauses satisfied}$$

$$E[Z] = \frac{1}{2} E[Z | x_1 = \text{true}] + \frac{1}{2} E[Z | x_1 = \text{false}]$$

$\Rightarrow$ either $E[Z | x_1 = \text{true}]$ or $E[Z | x_1 = \text{false}]$ is $\geq E[Z]$!

QuickSort,

Random divide and conquer

① pick random 'pivot'

② Divide A into two parts



Expected running time of QuickSort

For each $i, j \in [n]$ w/ $i < j$,

$X_{ij} = \begin{cases} 1 & \text{if } i\text{th smallest element is compared to } j\text{th smallest} \\ 0 & \text{otherwise} \end{cases}$

$\sum_{i,j} X_{ij}$ = total # comparisons made by the algorithm
 $\approx$ total running time

$E[\sum_{i,j} X_{ij}]$ = average running time

$$E[\sum_{i,j} X_{ij}] = \sum_{i,j} E[X_{ij}]$$

now analyze $E[X_{ij}]$ in isolation

$$E[X_{ij}] = \Pr[X_{ij} = 1] = \Pr[i\text{th element is compared to } j\text{th}]$$

i th and j th elem. selected before any element in between happens w/ probability $\frac{2}{j-i+1}$

$$E[X_{ij}] = \frac{2}{j-i+1}$$

$$E[\sum_{i,j} X_{ij}] = \sum_{i,j} \frac{2}{j-i+1}$$

$$\leq 2n \left(\frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right) \leq O(n \log n) \quad \text{with harmonic series}$$

Quickselect (HW)

today

Probability (& techniques)

Approximations via
linear programming
& randomized rounding

Algorithms

Max Sat

Set Cover

Covering
Integer Programs

Lecture 6 (today)

Probability (& techniques)

Approximations via
linear programming
& randomized rounding

Algorithms



Max Sat

Set Cover

Covering
Integer Programs

Randomized Rounding

Story of



Integer Programs (IPs)

min/max $\sum_{j=1}^n c_j x_j$

linear objective

over $x \in \mathbb{Z}^n$

integral variables

s.t. $A_1 x \leq b_1$
 $A_2 x = b_2$
 $A_3 x \geq b_3$

linear
constraints

for fixed vectors c, b_1, b_2, b_3 and matrices A_1, A_2, A_3

NP-Hard

Linear programs (LPs)

similar to IPs, except no integrality constraints

min/max $\sum_{j=1}^n c_j x_j$

linear objective

Over $x \in \cancel{\mathbb{Z}^n} \mathbb{R}^n$

integral variables

s.t. $A_1 x \leq b_1$

$$A_2 x = b_2$$

$$A_3 x \geq b_3$$

linear
constraints

for fixed vectors c, b_1, b_2, b_3 and matrices A_1, A_2, A_3

Linear Programs are in P (!!)

↑
(weakly polynomial)

1947 - Simplex algorithm
- Dantzig

(not polytime but practical)

1979 - Ellipsoid algorithm
- Khachiyan

polytime

1984 - Interior point algorithm
- Karmarkar

what's the difference?

Discrete
optimization

vs

Continuous
optimization

Integer
Programs

vs

Linear
Programs



NP-Hard



polynomial time!

LP-Based Approximations

① model as IP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{Over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1\end{array}$$




$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{Over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1\end{array}$$

② relax to LP



LP solver



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$



today

Probability (& techniques)

Approximations via
linear programming
& randomized rounding

Algorithms

Max Sat

Set Cover

Covering
Integer Programs

3-SAT

"conjunctive normal form"

"conjunctive normal form"
CNF formula

$$\varphi(x_1, \dots, x_n) = (x_1 \vee \overline{x_2} \vee x_3) \wedge (x_4 \vee \dots) \wedge \dots$$

m clauses, n variables

3 (distinct) variables per clause

The silliest APX algorithm

1. for $i = 1, \dots, n$

a. randomly set $x_i = \begin{cases} \text{TRUE} & \text{w/ prob. } .5 \\ \text{FALSE} & \text{w/ prob. } .5 \end{cases}$

We will:

① prove silly algo is $\frac{7}{8}$ -APX

② derandomize silly algo $\Rightarrow$ deterministic $7/8$ -APX

Even more surprising:

"probabilistically checkable proofs"

PCP Theorem

$(\frac{7}{8} + \epsilon)$ -APX is NP-Hard $\forall \epsilon > 0$

Approximating 3SAT: "Max 3-SAT"

$$\varphi(x_1, \dots, x_n) = (x_1 \vee \overline{x_2} \vee x_3) \wedge (x_4 \vee \dots) \wedge \dots$$


Goal: satisfy Max # clauses

Interesting even if φ is unsatisfiable

"d-Approximation algorithm" $d \in (0, 1)$

Algorithm satisfies $\underbrace{d \cdot \text{OPT}}$ clauses

Optimum (max) value



2 disjoint clauses

① prove silly algo is $\frac{7}{8}$ -APX

1. for $i=1, \dots, n$ The silliest APX algorithm
 a. randomly set $x_i = \begin{cases} \text{TRUE} & \text{w/ prob. } .5 \\ \text{FALSE} & \text{w/ prob. } .5 \end{cases}$

$$\mathcal{S}(x) = (\underbrace{x_1 \vee x_2 \vee \bar{x}_3}_{C_1}) \wedge (\underbrace{x_4 \vee \bar{x}_5 \vee x_6}_{C_2})$$

E_1 = event that C_1 satisfied

E_2 = $\sim$ C_2 $\sim$

$$\Pr[E_1, E_2] = \frac{7}{8} \cdot \frac{7}{8}$$

$$= \Pr[E_1] \cdot \Pr[E_2]$$

2 clauses w/
1 common variable

① prove silly algo is $\frac{7}{8}$ -APX

1. for $i=1, \dots, n$ The silliest APX algorithm
 a. randomly set $x_i = \begin{cases} \text{TRUE} & \text{w/ prob. } .5 \\ \text{FALSE} & \text{w/ prob. } .5 \end{cases}$

$$\mathcal{S}(x) = (\underbrace{x_1 \vee \bar{x}_2 \vee x_3}_{C_1}) \wedge (\underbrace{x_3 \vee x_4 \vee \bar{x}_5}_{C_2})$$

E_1 = event that C_1 satisfied

E_2 = $\sim$ C_2 $\sim$

$$\Pr[E_1, E_2] =$$


$$\underbrace{\Pr[E_1, E_2 | x_3 = T] \cdot \frac{1}{2}}_1 + \underbrace{\Pr[E_1, E_2 | x_3 = F] \cdot \frac{1}{2}}_{(\frac{3}{4})^2}$$

m clauses w/
common variables

① prove silly algo is $\frac{7}{8}$ -APX

1. for $i=1, \dots, n$ The silliest APX algorithm
 a. randomly set $x_i = \begin{cases} \text{TRUE} & \text{w/ prob. } .5 \\ \text{FALSE} & \text{w/ prob. } .5 \end{cases}$

$$\mathcal{S}(x) = (\underbrace{x_1 \vee \bar{x}_2 \vee x_3}_{C_1}) \wedge (\underbrace{x_3 \vee x_4 \vee \bar{x}_5}_{C_2}) \wedge (\underbrace{x_2 \vee \bar{x}_4 \vee x_6}_{C_3}) \wedge \dots$$



E_1 = event that C_1 satisfied

E_2 = $\sim$ C_2 $\sim$

E_3 = $\sim$ C_3 $\sim$

$\vdots$

general question

how to analyze a randomized mechanism
w/ a combinatorial explosion of possibilities?

Big idea: we only want the average

Linearity of Expectation

$$E[X+Y] = E[X] + E[Y] \quad (\text{ridiculously useful})$$

"average sum = sum of averages"

let $Y_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{otherwise} \end{cases}$

($i=1, \dots, m$)

$$E[Y_i] = \frac{7}{8} \quad (1 \text{ clause case})$$

$$E[\text{\# clauses satisfied}] = E\left[\sum_{i=1}^m Y_i\right] = \sum_{i=1}^m E[Y_i] = \frac{7}{8}m$$

(actually easy!)

Max-SAT

"conjunctive normal form"

CNF formula

$$\varphi(x_1, \dots, x_n) = (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_4 \vee \dots) \wedge \dots$$

m clauses, n variables

any # of variables per clause

goal: satisfy max # clauses

Approximation ratios for k-SAT vs max-SAT

k-sat

max-SAT

k oblivious APX

$$1 \quad 1/2$$

$$2 \quad 3/4$$

$$3 \quad 7/8$$

$$4 \quad 15/16$$

$$5 \quad 31/32$$

$\vdots$

$$k \quad 1 - 1/2^k$$

$$1/2$$

better?

LP-Based Approximations

① model as IP

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{Over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1\end{array}$$




$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{Over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1\end{array}$$

② relax to LP



LP solver



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$



IP for max-SAT

$$\max \sum_{i=1}^m z_i$$

$$\text{over } z_1, \dots, z_m \in \{0, 1\}$$

$$\gamma_1, \dots, \gamma_n \in \{0, 1\}$$

$$\sum_{i: x_i \in C_j} \gamma_i + \sum_{i: \bar{x}_i \in C_j} (1 - \gamma_i) \geq z_j \quad \forall j = 1, \dots, m$$

$$\text{clause } x_1 \vee \bar{x}_2 \vee x_3$$

$$z_1 \leq \gamma_1 + (1 - \gamma_2) + \gamma_3$$


LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

LP for max-SAT

$$\max \sum_{i=1}^m z_i$$

$$\text{over } y \in [0,1]^n, z \in [0,1]^m$$

$$\text{s.t. } \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i \quad \text{for each clause } C_i$$

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

e.g. $C_i = x_1 \vee \bar{x}_2 \vee x_3$

$\Rightarrow$ constraint

$$y_1 + (1 - y_2) + y_3 \geq z_1$$

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

IP for max-SAT

$$\max \sum_{i=1}^m z_i$$

over $y \in \{0,1\}^n, z \in \{0,1\}^m$

$$\text{s.t. } \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i$$

NP-Hard

LP for max-SAT

$$\max \sum_{i=1}^m z_i$$

over $y \in [0,1]^n, z \in [0,1]^m$

$$\text{s.t. } \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i$$

Solvable

$$\text{OPT}_{\text{IP}} \leq \text{OPT}_{\text{LP}}$$

Rounding LP solution

$$y \in [0,1]^n \rightsquigarrow x \in \{\text{TRUE}, \text{FALSE}\}$$

ideas?

$$\text{For } y_i, \quad x_i = \begin{cases} \text{TRUE} & \text{w/ prob. } y_i \\ \text{FALSE} & \text{w/ prob. } 1-y_i \end{cases}$$

$$x_i = \begin{cases} \text{TRUE} & \text{if } y_i \geq 1/2 \\ \text{FALSE} & \text{if } y_i < 1/2 \end{cases}$$

e.g., how to interpret $y_i = \frac{1}{2}$?

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

↑ rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$\begin{array}{ll} \max & \sum_{i=1}^m z_i \\ \text{over} & y \in [0,1]^n, z \in [0,1]^m \\ \text{s.t.} & \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1-y_j) \geq z_i \end{array}$$

for each clause C_i

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

Rounding LP solution

$$y \in [0, 1]^n \rightsquigarrow x \in \{\text{TRUE}, \text{FALSE}\}$$

For each variable x_j , independently, randomly set $x_j = \text{t}$ with probability y_j , and $x_j = \text{f}$ otherwise.

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

③ relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

↑ rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$\begin{array}{ll} \max & \sum_{i=1}^m z_i \\ \text{over} & y \in [0, 1]^n, z \in [0, 1]^m \\ \text{s.t.} & \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i \\ & \text{for each clause } C_i \end{array}$$

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

For each variable x_j , independently, randomly set $x_j = \text{t}$ with probability y_j , and $x_j = \text{f}$ otherwise.

claim: each clause C_i is satisfied
w/ probability $(1 - 1/e) z_i$

proof:


$$P[\text{don't satisfy } C_i] =$$

$$= \prod_{j: x_j \in C_i} P[x_j = \text{f}] \prod_{j: \bar{x}_j \in C_i} P[x_j = \text{t}]$$

$$= \prod_{j: x_j \in C_i} (1 - y_j) \prod_{j: \bar{x}_j \in C_i} y_j$$

$$\leq \prod_{j: x_j \in C_i} e^{-y_j} \prod_{j: \bar{x}_j \in C_i} e^{-(1-y_j)}$$

$$= e^{-(\sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1-y_j))} \leq e^{-z_i}$$



$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$$1 + x \leq e^x$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP




approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$e^{-z_i}$$

$$f(x) = e^{-x}$$



$$e^{-z_i} \leq \underbrace{(1-z_i)}_{1} e^{\underbrace{0}_{1/e}} + z_i e^{-1} = \underline{1 - (1 - 1/e)z_i}$$

$$P[\text{don't satisfy } C_i] \leq 1 - (1 - 1/e)z_i$$

$$P[\text{satisfy } C_i] \geq (1 - 1/e)z_i$$

For each variable x_j , independently, randomly set $x_j = t$ with probability y_j , and $x_j = f$ otherwise.

claim: each clause C_i is satisfied
w/ probability $(1 - 1/e) z_i$

claim:

$$E[\# \text{ sat. clauses}] \geq (1 - 1/e) \text{OPT}$$

$$\hookrightarrow \geq \sum_i P[\text{satisfy } C_i]$$

$$\geq (1 - 1/e) \sum_i z_i$$

$$= (1 - 1/e) \text{OPT}_{LP} \geq (1 - 1/e) \text{OPT}_{SAT}$$

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$\max \sum_{i=1}^m z_i$$

$$\text{over } y \in [0, 1]^n, z \in [0, 1]^m$$

$$\text{s.t. } \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i$$

for each clause C_i

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

For each variable x_j , independently, randomly set $x_j = \text{t}$ with probability y_j , and $x_j = \text{f}$ otherwise.

$$E[\# \text{ sat. clauses}] \geq (1 - 1/e) \text{OPT}$$

Better?

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

③ relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

↑ rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$\begin{array}{ll} \max & \sum_{i=1}^m z_i \\ \text{over} & y \in [0,1]^n, z \in [0,1]^m \\ \text{s.t.} & \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i \end{array}$$

for each clause C_i

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

For each variable x_j , independently, randomly set $x_j = t$ with probability y_j , and $x_j = f$ otherwise.

claim: each clause C_i is satisfied
w/ probability $(1 - 1/e) z_i$
.68

what if C_i has 1 variable?

$$C_i = x_1 \quad P[\text{sat. } C_i] = y_1 = z_i$$

~ 2 variables?

$$C_i = x_1 \vee x_2$$

$$P[\text{don't sat } C_i] = (1 - y_1)(1 - y_2)$$

$$\leq \left(\frac{(1 - y_1) + (1 - y_2)}{2} \right)^2$$

$$\leq \left(1 - \frac{z_i}{2} \right)^2 \leq (1 - z_i) \left(1 - \frac{0}{2} \right)^2 + z_i \left(1 - \frac{1}{2} \right)^2$$

$$f(x) = \left(1 - \frac{x}{2} \right)^2 \quad y_1 + y_2 \geq z_i$$


LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$
↑ rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

$$\max \sum_{i=1}^m z_i$$

$$\text{over } y \in [0, 1]^n, z \in [0, 1]^m$$

$$\text{s.t. } \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i$$

for each clause C_i

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$


$$\sqrt{ab} \leq \frac{a+b}{2}$$

$$= 1 - z_i + \frac{z_i}{4} = 1 - \frac{3}{4}z_i$$

$$P[\text{sat } C_i] \geq \frac{3}{4}z_i$$

For each variable x_j , independently, randomly set $x_j = \text{t}$ with probability y_j , and $x_j = \text{f}$ otherwise.

$$E[\# \text{ sat. clauses}] \geq (1 - 1/e) \text{OPT}$$



k	oblivious	LP	average
1	1/2	z_i	$\geq (3/4)z_i$
2	3/4	$(3/4)z_i$	$\geq (3/4)z_i$
3	7/8	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
4	15/16	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
5	31/32	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
$\vdots$			
k	$1 - 1/2^k$	$(1 - 1/e)z_i$	$\geq (3/4)z_i$

$$\begin{array}{ll} \max & \sum_{i=1}^m z_i \\ \text{over} & y \in [0,1]^n, z \in [0,1]^m \\ \text{s.t.} & \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i \\ & \text{for each clause } C_i \end{array}$$

$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$

$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$

Can we get $\frac{3}{4}$ -APX?

- ① Flip a coin
- ② If heads, do oblivious rounding
- ③ If tails, do LP rounding

For each variable x_j , independently, randomly set $x_j = t$ with probability y_j , and $x_j = f$ otherwise.


claim: each clause C_i is satisfied
w/ probability $\frac{3}{4} z_i$

say C_i has k clauses

$$P[\text{sat } C_i] = P[\text{sat } C_i | \text{heads}] \cdot \frac{1}{2} + P[\text{sat } C_i | \text{Tails}] \cdot \frac{1}{2}$$

$$\geq (1 - \frac{1}{2^k}) \frac{1}{2} + \left\{ \frac{1}{3/4} \quad \begin{matrix} k=1 \\ k=2 \end{matrix} \right\} \frac{1}{2}$$

$$\geq \frac{3}{4}$$



$$\begin{array}{ll} \max & \sum_{i=1}^m z_i \\ \text{over} & y \in [0,1]^n, z \in [0,1]^m \\ \text{s.t.} & \sum_{j: x_j \in C_i} y_j + \sum_{j: \bar{x}_j \in C_i} (1 - y_j) \geq z_i \end{array}$$

for each clause C_i

$$z_i = \begin{cases} 1 & \text{if } i\text{th clause satisfied} \\ 0 & \text{if not} \end{cases}$$

$$y_j = \begin{cases} 1 & \text{if } x_j = \text{TRUE} \\ 0 & \text{if } x_j = \text{FALSE} \end{cases}$$

k	oblivious	LP	average
1	1/2	z_i	$\geq (3/4)z_i$
2	3/4	$(3/4)z_i$	$\geq (3/4)z_i$
3	7/8	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
4	15/16	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
5	31/32	$(1 - 1/e)z_i$	$\geq (3/4)z_i$
⋮			
k	$1 - 1/2^k$	$(1 - 1/e)z_i$	$\geq (3/4)z_i$

today

Probability (& techniques)

Approximations via
linear programming
& randomized rounding

Algorithms

Max Sat


Set Cover

Covering
Integer Programs

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$



Goal: select k sets $S_{e_1}, S_{e_2}, \dots, S_{e_k}$,
minimizing k ,

s.t. $S_{e_1} \cup S_{e_2} \cup \dots \cup S_{e_k} = [m]$


NP-Hard

$$\varphi(x_1, \dots, x_n) =$$



$$(x_1 \vee \bar{x}_2)$$

$$\wedge (x_2 \vee x_3 \vee \bar{x}_4)$$

$$\wedge m$$



yes, no



[pause here and try
to think of a
reduction from 3-SAT]


$$f(x) = (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_2 \vee x_4 \vee x_5) \wedge (\bar{x}_1 \vee x_3 \vee \bar{x}_5) \wedge \dots \left(\begin{array}{l} n \text{ variables,} \\ m \text{ clauses} \end{array} \right)$$

elements
per
clause

sets =
assignments

elements
per variable

IP for set cover



Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$

minimizing k ,

s.t. $S_{e_1} \cup \dots \cup S_{e_k} = [m]$

IP for set cover

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in \{0, 1\}^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



optimal LP solution $x \in \mathbb{R}^n$

↑ rounding

approximate integral solution $z \in \mathbb{Z}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$

minimizing k ,

s.t. $S_{e_1} \cup \dots \cup S_{e_k} = [m]$

LP for set cover

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$

minimizing k ,

s.t. $S_{e_1} \cup \dots \cup S_{e_k} = [m]$

IP for set cover

$$\min \sum_{j=1}^n x_j$$

over $x \in \{0,1\}^n$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \\ \forall i \in [m]$$

NP-Hard

LP for set cover

$$\min \sum_{j=1}^n x_j$$

over $x \in [0,1]^n$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \\ \forall i \in [m]$$


Polytime solvable

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

Ideas?

How to interpret $x_i = .5$?



Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

For each set S_j , independently, take S_j with probability x_j .

$$E[|F|] = \sum_j x_j = \text{OPT} \quad (\text{good!})$$

Fix a point i .

$$E[\text{sets in } F \text{ covering } i]$$

$$P[F \text{ not cover } i]$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

For each set S_j , independently, take S_j with probability x_j .

$$E[|F|] = \sum_j x_j = \text{OPT} \quad (\text{good!})$$

Fix a point i .

$$E[\text{sets in } F \text{ covering } i] = \sum_{j: i \in S_j} x_j \geq 1$$

$$P[F \text{ not cover } i] = \prod_{j: i \in S_j} (1 - x_j) \leq \prod_{j: i \in S_j} e^{-x_j}$$

$$= e^{-\sum_{j: i \in S_j} x_j} \leq e^{-1}$$

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

over $x \in [0,1]^n$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

For each set S_j , independently, take S_j with probability x_j .

$$E[|F|] = \sum_j x_j = \text{OPT} \quad (\text{good!})$$

$$P[F \text{ not cover } i] = e^{-1}$$

so F covers each point w/
constant probability

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding
optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

2. For each set S_j independently, take S_j with probability $\min\{1, \alpha x_j\}$.

for $d > 1$ TBD.

$$E[|F|] =$$

$$P[F \text{ not cover } i]$$

$$P[F \text{ not set cover}]$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

2. For each set S_j independently, take S_j with probability $\min\{1, \alpha x_j\}$.

for $d > 1$ TBD.

$$E[|F|] = d \sum_j x_j = d \text{OPT}$$

$$P[F \text{ not cover } i] = \prod_{S_j \ni i} (1 - d x_j) \leq e^{-d \sum_{S_j \ni i} x_j} \leq e^{-d}$$

$$P[F \text{ not set cover}] \leq \sum_i P[F \text{ not cover } i] \leq m \cdot e^{-d} \leq \frac{1}{m^{\alpha}} \text{ for } d = O(\log m)$$


LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_j x \leq b_j \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_j x \leq b_j \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$
minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

Rounding LP solution

$$x \in [0,1]^n \rightsquigarrow F \subseteq \{S_1, \dots, S_n\}$$

2. For each set S_j independently, take S_j with probability $\min\{1, \alpha x_j\}$.

for $d > 1$ TBD.

$$E[|F|] = d \sum_j x_j = d \text{OPT}$$

$$P[F \text{ not set cover}] \leq \frac{1}{m^{\alpha d}}$$

$$\text{for } d = O(\log m)$$

$$\Rightarrow O(\log m)\text{-APX}$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

Minimum Set Cover

m points $[m] \equiv \{1, \dots, m\}$

n sets $S_1, \dots, S_n \subseteq [m]$

Goal: select k sets $S_{e_1}, \dots, S_{e_k}$

minimizing k ,

$$\text{s.t. } S_{e_1} \cup \dots \cup S_{e_k} = [m]$$

$$\min \sum_{j=1}^n x_j$$

$$\text{over } x \in [0,1]^n$$

$$\text{s.t. } \sum_{j: i \in S_j} x_j \geq 1 \quad \forall i \in [m]$$

$$x_j = \begin{cases} 1 & \text{if } S_j \text{ in set cover,} \\ 0 & \text{otherwise} \end{cases}$$

today

Probability (& techniques)

Approximations via
linear programming
& randomized rounding

Algorithms

Max Sat

Set Cover

Covering
Integer Programs

Covering IPs

$$\min \sum_j c_j x_j$$

$$\text{over } x \in \mathbb{Z}_{\geq 0}^n$$

$$\text{s.t. } \sum_{j=1}^n A_{ij} x_j \geq b_i \text{ for } i \in [m]$$

$$\text{where } A \in \mathbb{R}_{\geq 0}^{m \times n}, \quad b \in \mathbb{R}_{> 0}^m, \quad c \in \mathbb{R}_{> 0}^n$$

"demands" "costs"

WLOG $A_{ij} \leq b_i \quad \forall i, j$ (why?)


Set cover is special case w/ $A_{ij} \in \{0, 1\}$, $b_i = 1$, $c_j = 1$

LP relaxation

$$\min \sum_j c_j x_j$$

$$\text{over } x \in \mathbb{Z}_{\geq 0}^n \quad \mathbb{R}_{\geq 0}^n$$

$$\text{s.t. } \sum_{j=1}^n A_{ij} x_j \geq b_i \quad \text{for } i \in [m]$$



Rounding LP solution

$$x \in \mathbb{R}_{\geq 0}^n \rightsquigarrow z \in \mathbb{Z}_{\geq 0}^n$$

Ideas?

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$



$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



optimal LP solution
 $x \in \mathbb{R}^n$

approximate integral solution
 $z \in \mathbb{Z}^n$
↑ rounding

LP relaxation

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{R}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \end{array}$$

Covering IPs

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{Z}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \\ & A_{ij} \leq b_i \quad \forall i, j \end{array}$$

Rounding LP solution

$$x \in \mathbb{R}_{\geq 0}^n \rightsquigarrow z \in \mathbb{Z}_{\geq 0}^n$$

For each $j \in [n]$, independently, let

$$z_j = \begin{cases} \lceil \alpha x_j \rceil & \text{with probability } \alpha x_j - \lfloor \alpha x_j \rfloor \\ \lfloor \alpha x_j \rfloor & \text{otherwise.} \end{cases}$$

$$E[z_j] = \alpha x_j$$

$$E\left[\sum_j c_j z_j\right] = \alpha \sum_j c_j x_j = \alpha \text{OPT}$$

LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

↓

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

↑ rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

LP relaxation

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{R}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \end{array}$$

for $i \in [m]$

Covering IPs

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{Z}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \end{array}$$

for $i \in [m]$

$A_{ij} \leq b_i \quad \forall i, j$

For each $j \in [n]$, independently, let

$$z_j = \begin{cases} \lceil \alpha x_j \rceil & \text{with probability } \alpha x_j - \lfloor \alpha x_j \rfloor \\ \lfloor \alpha x_j \rfloor & \text{otherwise.} \end{cases}$$

$$P[\sum_j A_{ij} z_j \leq b_i]$$


LP-Based Approximations

① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP



LP solver

optimal LP solution $x \in \mathbb{R}^n$

approximate integral solution $z \in \mathbb{Z}^n$

rounding

LP relaxation

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{R}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \end{array}$$

Covering IPs

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{Z}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \\ & A_{ij} \leq b_i \quad \forall i, j \end{array}$$

For each $j \in [n]$, independently, let

$$z_j = \begin{cases} \lceil \alpha x_j \rceil & \text{with probability } \alpha x_j - \lfloor \alpha x_j \rfloor \\ \lfloor \alpha x_j \rfloor & \text{otherwise.} \end{cases}$$

$$P[\sum_j A_{ij} z_j \leq b_i]$$

LP-Based Approximations

① model as IP

$$\begin{aligned} \max \quad & \sum_{j=1}^n c_j x_j \\ \text{over } & x \in \mathbb{Z}^n \\ \text{s.t. } & A_1 x \leq b_1 \end{aligned}$$

$$\begin{aligned} \max \quad & \sum_{j=1}^n c_j x_j \\ \text{over } & x \in \mathbb{R}^n \\ \text{s.t. } & A_1 x \leq b_1 \end{aligned}$$

② relax to LP



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding
optimal
LP solution
 $x \in \mathbb{R}^n$

LP relaxation

$$\begin{aligned} \min \quad & \sum_j c_j x_j \\ \text{over } & x \in \mathbb{R}_{\geq 0}^n \\ \text{s.t. } & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \end{aligned}$$

Covering IPs

$$\begin{aligned} \min \quad & \sum_j c_j x_j \\ \text{over } & x \in \mathbb{Z}_{\geq 0}^n \\ \text{s.t. } & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \\ & A_{ij} \leq b_i \quad \forall i, j \end{aligned}$$

Multiplicative Chernoff bound

let $X_1, \dots, X_n \in [0, 1]$ be independent

$$\mu = E[X_1 + \dots + X_n] \in (0, 1)$$

$$P[X_1 + \dots + X_n \leq (1 - \epsilon)\mu] \leq e^{-\epsilon^2 \mu / 2}$$

For each $j \in [n]$, independently, let

$$z_j = \begin{cases} \lceil \alpha x_j \rceil & \text{with probability } \alpha x_j - \lfloor \alpha x_j \rfloor \\ \lfloor \alpha x_j \rfloor & \text{otherwise.} \end{cases}$$

LP-Based Approximations


① model as IP

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{Z}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

$$\begin{array}{ll} \max & \sum_{j=1}^n c_j x_j \\ \text{over} & x \in \mathbb{R}^n \\ \text{s.t.} & A_1 x \leq b_1 \end{array}$$

② relax to LP

LP solver



approximate
integral solution
 $z \in \mathbb{Z}^n$

rounding

optimal
LP solution
 $x \in \mathbb{R}^n$

LP relaxation

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{R}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \end{array}$$

Covering IPs

$$\begin{array}{ll} \min & \sum_j c_j x_j \\ \text{over} & x \in \mathbb{Z}_{\geq 0}^n \\ \text{s.t.} & \sum_{j=1}^n A_{ij} x_j \geq b_i \\ & \text{for } i \in [m] \\ & A_{ij} \leq b_i \quad \forall i, j \end{array}$$

Multiplicative Chernoff bound
let $X_1, \dots, X_n \in [0, 1]$ be independent

$$\mu = \mathbb{E}[X_1 + \dots + X_n] \in (0, 1)$$

$$P[X_1 + \dots + X_n \leq (1 - \epsilon)\mu] \leq e^{-\epsilon^2 \mu / 2}$$

Multiplicative Chernoff bound

let $X_1, \dots, X_n \in [0, 1]$ be independent

$$\mu = E[X_1 + \dots + X_n]$$

$$\epsilon \in (0, 1)$$

$$\bullet \quad P[X_1 + \dots + X_n \geq (1 + \epsilon)\mu] \leq e^{-\epsilon^2 \mu / 3}$$

$$\bullet \quad P[X_1 + \dots + X_n \leq (1 - \epsilon)\mu] \leq e^{-\epsilon^2 \mu / 2}$$

(proof at end if time permits)